

ON LINEAR REPRESENTATIONS OF SOME EXTENSIONS

V. G. BARDAKOV, O. V. BRYUKHANOV

Abstract. Yu. I. Merzljakov developed a method of splittable coordinates which helps to verify the linearity of some groups, he established some fundamental results using this method.

In this paper we use the method of splittable coordinates and find some sufficient condition under which the semi-direct product of two linear groups is linear. As consequence we get linearity of some HNN-extensions of a free group, linearity of the holomorph of the braid group B_n , $n \geq 2$, and free group F_2 and linearity of some Artin groups. In all cases we construct faithful linear representations in the explicit form.

AMS Classification: 20F28; 20F36.

Keywords: linear group, faithful linear representation, HNN-extension, semi-direct product.

A. I. Malcev [1] formulated the problem of characterization of abstract groups which has faithful linear representation over some field (we will said that these groups are linear groups). A. I. Malcev found the necessary and sufficient conditions under which the abelian group has a faithful linear representation. It is known some results on linear representations of nilpotent, polycyclic, solvable and some other groups (see the survey of Yu. I. Merzljakov [2]). A. Lubotzky [3] found a characterization of abstract groups which has faithful linear representation over field of complex numbers $\mathbb{C}$. But this result is hard to use for the concrete groups.

Another direction of investigation is the following problem: if a group G is constructed from linear groups with the help of some group construction (free product, free product with amalgamation, extension, HNN-extension, wreath product and so on), then for which conditions group G is linear? In this direction V. L. Nisnevic [4] proved that a free product of two linear groups is a linear group. B. A. F. Wehrfritz [5] found some sufficient conditions under which a free product with amalgamation of linear groups is a linear group. Yu. E. Vapne [6] studied a wreath product of linear groups. Also he proved sufficient conditions of linearity of some nilpotent product [7]. O. V. Bryukhanov [8] found the necessary and sufficient conditions under which the nilpotent product $A(n)B$, $n \geq 2$, of finitely generated linear groups A and B has a faithful linear representation over fields of zero and positive characteristics.

But the question of existence of faithful linear representations for group extensions and HNN-extensions is still open. We have few results in this area. In particular, it is not known is it true that a semi-direct product of two finitely generated linear groups is linear. R. T. Volvachev [9] solved the question about faithful linear representations of HNN-extensions a cyclic group. By well-known result of A. I. Malcev [1] every semi-direct product of residually finite groups is residually finite. On the other hand it

is known that an extension of a linear group by a linear group can be non-linear. As an example we can consider a finitely generated free solvable group of class 3. This group is not linear but is the extension of free metabelian group of infinite rank by finitely generated abelian group.

Yu. I. Merzljakov [10], [11] developed a method (the method of splittable coordinates) which helps to prove linearity of some groups. Yu. I. Merzljakov established some fundamental results using this method.

In this paper, we will use the method of splittable coordinates and find some sufficient condition under which the semi-direct product of two linear groups is linear (see Theorem 1). Also we establish linearity of $\text{Hol}(B_n)$, $n \geq 3$ and $\text{Hol}(F_2)$ over field of characteristic zero from these follows linearity over field of characteristic zero any splittable extension of braid group B_n , $n \geq 3$, and free group F_2 by linear group.

In Section 2 we consider HNN-extensions

$$F_\varphi(X) = \langle X, t \mid t^{-1}xt = \varphi(t), \quad x \in X \rangle$$

of a free group $F(X)$ with a virtually inner automorphism φ of $F(X)$ (recall that an automorphism φ is called virtually inner if φ^m is inner automorphism of $F(X)$ for some positive integer m). Note that these groups are studied in the papers [12], [13]. We will construct a faithful linear representation of $F_\varphi(X)$ (see Theorem 2). Also we will find the center of $F_\varphi(X)$.

In the last Section we study 2-generated Artin groups and construct a faithful linear representations over ring of rational fractions $\mathbb{Q}_p$ with denominators equal to powers of prime number p . C. C. Squier [14] established the linearity of 2-generated Artin groups but he constructed representation over field with transcendental (over $\mathbb{Q}$) numbers.

Acknowledgements. We would like to thank M. V. Neshchadim for his remarks on the first draft of this paper. Special thanks goes to the participants of the seminar “Evariste Galois” at Novosibirsk State University for their kind attention to our work. Authors were supported by RFBR (grant 02-01-01118).

§ 1. Method of splittable coordinates

We remind that some homomorphism $\rho : G \longrightarrow \text{GL}_n(K)$ of group G into general linear group $\text{GL}_n(K)$ over field K is called *the linear representation*. This representation ρ is called *faithful* if the kernel $\text{Ker}(\rho)$ is trivial. A group is called *linear* if it has some faithful linear representation. Yu. I. Merzljakov [10], [11] developed a method (the method of splittable coordinates) which helps to prove linearity of some groups. The heart of this method is to assign a faithful action of group on finite dimension vector space of algebra of functions from group G to field K . Let $A(G, K)$ be the algebra of all K -valuable functions on G . If $f \in A(G, K)$, $x \in G$ then we define the action of x by the rule

$$f^x(y) = f(xy), \quad y \in G.$$

Since $f^{gh} = (f^g)^h$ and $f^e = f$, where e is the unit of G , $g, h \in G$ then we get the action of G on the algebra $A(G, K)$.

We will call any subset of functions $\{t_i\} \subseteq A(G, K)$, $i \in I$, *K-valuable coordinates* on G , if for any $x \in G \setminus \{e\}$ there is $t_j \in \{t_i\}$ such that $t_j(x) \neq t_j(e)$. We will say that the K -valuable coordinates $t_1, \dots, t_d$ on G , are *splittable coordinates* if the following equality are true:

$$t_\alpha(xy) = \sum_{\beta=1}^l f_{\alpha\beta}(x)h_\beta(y), \quad x, y \in G, \quad 1 \leq \alpha \leq d,$$

where $f_{\alpha\beta}, h_\beta$ are some K -valuable functions on G .

In the case when the K -valuable splittable coordinates $t_1, \dots, t_d$ are defined on G we can see that a vector space V generated by G -orbit of the set of coordinate functions $\{t_\alpha\}$ has finite dimension because it lies in linear K -envelope of the finite set $\{h_\beta\}$. Let $v_1, \dots, v_n$ be a basis of V over K . Then for every $x \in G$ we have

$$v_i^x = \sum_{j=1}^n \rho_{ij}(x)v_j, \quad \rho_{ij}(x) \in K.$$

The linear representation ρ which is defined by the rule $x \mapsto (\rho_{ij}(x))$, $x \in G$ is faithful. Namely, if the identical linear transformation of V corresponding to some element x from G then $t_\alpha^x = t_\alpha$, where $1 \leq \alpha \leq d$. In this case $t_\alpha(x) = t_\alpha^x(e) = t_\alpha(e)$ if $1 \leq \alpha \leq d$ and hence $x = e$. The following lemma describes some properties of this representation (see [10], [11]).

Lemma 1 (on splittable coordinates). *Let K be a field and G be a group with K -valuable coordinates $t_1, \dots, t_d$ and*

$$t_\alpha(xy) = \sum_{\beta=1}^l f_{\alpha\beta}(x)h_\beta(y), \quad x, y \in G, \quad 1 \leq \alpha \leq d,$$

where $f_{\alpha\beta}, h_\beta$ are functions with the values in K . There exists the isomorphism $\rho : G \rightarrow \text{GL}_n(K)$ such that: 1) ρ^{-1} is linear on G^p , 2) if the function $f_{\alpha\beta}$ is polynomial on G (in coordinates t_α) and h_β is polynomial on subset $G_1 \subset G$ then ρ is polynomial on G_1 .

We will say that the map $f : X \times Y \mapsto \text{Mat}_n(K)$ is *splittable over K of length l* if for all its components f_{ij} the following equality are true

$$f_{ij}(x, y) = \sum_{\nu=1}^l \varphi_{\nu ij}(x)\psi_{\nu ij}(y), \quad x \in X, y \in Y, \quad 1 \leq i, j \leq n,$$

where $\varphi_{\nu ij}, \psi_{\nu ij}$ are the functions with the values in K . This yields that the coordinates $\{t_\alpha\}$ are splittable if the functions $(x, y) \mapsto t_\alpha(xy)$ are splittable.

The following statement was formulated in [11] as an exercise. We give the proof of this statement.

Lemma 2. Let G be an extension of a group A by a group B with the factors $f(x, y) \in A$, $x, y \in B$, and automorphisms $\hat{x} \in \text{Aut}(A)$, $x \in B$. If A and B are linear over the field K and the maps

$$(x, y) \mapsto f(x, y), \quad (a, x) \mapsto a^{\hat{x}}$$

are splittable then the group G is linear over K .

Proof. Since G is an extension of A by B then for every $g \in G$ there exists a single pair (b, a) , $b \in B, a \in A$ and the product of these pairs is defined by

$$g_1 g_2 = (b_1, a_1) \cdot (b_2, a_2) = (b_1 b_2, f(b_1, b_2) a_1^{\hat{b}_2} a_2).$$

Since A and B are linear then there exist the faithful linear representations:

$$\rho_A : A \longrightarrow \text{GL}_m(K), \quad \rho_B : B \longrightarrow \text{GL}_n(K).$$

We take the following functions as coordinates on A and B :

$$t_{ij}^{(A)}(a) = (\rho_A(a))_{ij}, \quad a \in A, \quad 1 \leq i, j \leq m, \quad t_{ij}^{(B)}(b) = (\rho_B(b))_{ij}, \quad b \in B, \quad 1 \leq i, j \leq n.$$

It is evident that the set of functions

$$\{T_{ij}^{(A)} \mid 1 \leq i, j \leq m\} \cup \{T_{ij}^{(B)} \mid 1 \leq i, j \leq n\},$$

where

$$T_{ij}^{(A)}((b, a)) = t_{ij}^{(A)}(a), \quad T_{ij}^{(B)}((b, a)) = t_{ij}^{(B)}(b),$$

are coordinates on G .

Let us show that these coordinates $T_{ij}^{(A)}$, $T_{ij}^{(B)}$ are splittable. For this we calculate

$$T_{ij}^{(A)}(g_1 g_2) = t_{ij}^{(A)}(f(b_1, b_2) a_1^{\hat{b}_2} a_2) = \sum_{k,l} t_{ik}^{(A)}(f(b_1, b_2)) t_{kl}^{(A)}(a_1^{\hat{b}_2}) t_{lj}^{(A)}(a_2).$$

Since the maps

$$(x, y) \mapsto f(x, y), \quad (a, x) \mapsto a^{\hat{x}}$$

are splittable then

$$t_{ik}^{(A)}(f(b_1, b_2)) = \sum_{\nu} \varphi_{\nu ik}(b_1) \psi_{\nu ik}(b_2), \quad t_{kl}^{(A)}(a_1^{\hat{b}_2}) = \sum_{\mu} \bar{\varphi}_{\mu kl}(a_1) \bar{\psi}_{\mu kl}(b_2).$$

Therefore,

$$\begin{aligned} & \sum_{k,l} t_{ik}^{(A)}(f(b_1, b_2)) t_{kl}^{(A)}(a_1^{\hat{b}_2}) t_{lj}^{(A)}(a_2) = \\ &= \sum_{k,l} \left(\sum_{\nu} \varphi_{\nu ik}(b_1) \psi_{\nu ik}(b_2) \right) \left(\sum_{\mu} \bar{\varphi}_{\mu kl}(a_1) \bar{\psi}_{\mu kl}(b_2) \right) t_{lj}^{(A)}(a_2) = \\ &= \sum_{k,l,\mu,\nu} (\varphi_{\nu ik}(b_1) \bar{\varphi}_{\mu kl}(a_1)) (\psi_{\nu ik}(b_2) \bar{\psi}_{\mu kl}(b_2)) t_{lj}^{(A)}(a_2). \end{aligned}$$

Let

$$\Phi_\alpha(g_1) = \varphi_{\nu ik}(b_1)\overline{\varphi}_{\mu kl}(b_1), \quad \Psi_{\alpha j}(g_2) = \psi_{\nu ik}(b_2)\overline{\psi}_{\mu kl}(b_2)t_{lj}^{(A)}(a_2), \quad \alpha = (i, k, l, \mu, \nu).$$

then we have

$$T_{ij}^{(A)}(g_1 g_2) = \sum_{\alpha} \Phi_{\alpha}(g_1) \Psi_{\alpha j}(g_2).$$

The set of functions $\{\Psi_{\alpha j}\}$ is finite since the indexes i, j, k, l, μ, ν may take only finite number of values. This implies that the coordinates $T_{ij}^{(A)}$ are splittable.

For the functions $T_{ij}^{(B)}$ we have

$$T_{ij}^{(B)}(g_1 g_2) = t_{ij}^{(B)}(b_1 b_2) = \sum_k t_{ik}^{(B)}(b_1) t_{kj}^{(B)}(b_2) = \sum_k T_{ik}^{(B)}(g_1) T_{kj}^{(B)}(g_2).$$

Therefore the coordinates $T_{ij}^{(A)}, T_{ij}^{(B)}$ are splittable and linearity of G follows from Lemma 1.

Now let us prove the following theorem

Theorem 1. *Let K be a field of arbitrary characteristic, $\Phi \leq \text{GL}_m(K)$, $G \leq \text{GL}_n(K)$. If for every $\varphi \in \Phi$ there is $g_{\varphi} \in \text{GL}_n(K)$ such that $\varphi^{-1}g\varphi = g_{\varphi}^{-1}gg_{\varphi}$ for all $g \in G$ then the group $\Phi \rtimes G$ has faithful linear representation by matrices of size $\leq m^2 + n^4$ over the field K .*

Proof. For every element from $\Phi \rtimes G$ there is only one pair (φ, g) , $\varphi \in \Phi$, $g \in G$ which represents this element. Then on the group $\Phi \rtimes G$ we can define the coordinate functions $T_{ij}^{(1)}$, $1 \leq i, j \leq n^2$, $T_{pq}^{(2)}$, $1 \leq p, q \leq n$, by setting

$$T_{ij}^{(1)}(\varphi, g) = t_{ij}^{(1)}(\varphi), \quad T_{pq}^{(2)}(\varphi, g) = t_{pq}^{(2)}(g),$$

where $t_{ij}^{(1)}$, $t_{pq}^{(2)}$ are the components of faithful linear representations of Φ and $\text{GL}_n(K)$ correspondingly, i. e. the coordinate functions on Φ and on $\text{GL}_n(K)$.

For every automorphism $\varphi \in \Phi$ we take element $g_{\varphi} \in \text{GL}_n(K)$ such that $g^{\varphi} = g_{\varphi}^{-1}gg_{\varphi}$ for all $g \in G$. We define this function by $\tau : \varphi \mapsto g_{\varphi}$. Since $(\varphi_1, g_1)(\varphi_2, g_2) = (\varphi_1\varphi_2, g_1^{\varphi_2}g_2)$ then

$$\begin{aligned} T_{ij}^{(1)}((\varphi_1, g_1)(\varphi_2, g_2)) &= T_{ij}^{(1)}(\varphi_1\varphi_2, g_1^{\varphi_2}g_2) = t_{ij}^{(1)}(\varphi_1\varphi_2) = \sum_{k=1}^{n^2} t_{ik}^{(1)}(\varphi_1)t_{kj}^{(1)}(\varphi_2) = \\ &= \sum_{k=1}^{n^2} T_{ik}^{(1)}(\varphi_1, g_1)T_{kj}^{(1)}(\varphi_2, g_2), \\ T_{pq}^{(2)}((\varphi_1, g_1)(\varphi_2, g_2)) &= T_{pq}^{(2)}(\varphi_1\varphi_2, g_1^{\varphi_2}g_2) = t_{pq}^{(2)}(g_1^{\varphi_2}g_2) = t_{pq}^{(2)}(g_{\varphi_2}^{-1}g_1g_{\varphi_2}g_2) = \\ &= \sum_{k_1, k_2, k_3=1}^n t_{pk_1}^{(2)}(g_{\varphi_2}^{-1})t_{k_1k_2}^{(2)}(g_1)t_{k_2k_3}^{(2)}(g_{\varphi_2})t_{k_3q}^{(2)}(g_2) = \\ &= \sum_{k_1, k_2, k_3=1}^n t_{k_1k_2}^{(2)}(g_1)t_{pk_1}^{(2)}(g_{\varphi_2}^{-1})t_{k_2k_3}^{(2)}(g_{\varphi_2})t_{k_3q}^{(2)}(g_2) = \end{aligned}$$

$$\begin{aligned}
&= \sum_{k_1, k_2=1}^n t_{k_1 k_2}^{(2)}(g_1) \left(\sum_{k_3=1}^n t_{p k_1}^{(2)}(g_{\varphi_2}^{-1}) t_{k_2 k_3}^{(2)}(g_{\varphi_2}) t_{k_3 q}^{(2)}(g_2) \right) = \\
&= \sum_{k_1, k_2=1}^n T_{k_1 k_2}^{(2)}(\varphi_1, g_1) H_{p k_1 k_2 q}(\varphi_2, g_2),
\end{aligned}$$

where

$$H_{p k_1 k_2 q}(\varphi_2, g_2) = \sum_{k_3=1}^n t_{p k_1}^{(2)}(g_{\varphi_2}^{-1}) t_{k_2 k_3}^{(2)}(g_{\varphi_2}) t_{k_3 q}^{(2)}(g_2).$$

Since the set of functions $\{T_{ij}^{(1)}, H_{p k_1 k_2 q}\}$ is finite and it has $m^2 + n^4$ functions then by Lemma 1 the group $\Phi \rtimes G$ has a faithful linear representation by the matrices of size $\leq m^2 + n^4$ over the field K . This completes the proof.

From this theorem easily follows

Corollary 1. *Let K be a field of arbitrary characteristic and $G \leq \text{GL}_n(K)$. Then the group $\text{Int}(G)G$ embeds in $\text{GL}_{2n^4}(K)$.*

Corollary 2. *The holomorph $\text{Hol}(B_n)$ of the braid group B_n , $n \geq 2$, has a faithful linear representation over a field of characteristic zero.*

Proof. As was proved in [16] $|\text{Aut}(B_n) : \text{Int}(B_n)| = 2$. Hence, $|\text{Hol}(B_n) : \text{Int}(B_n)B_n| = 2$. Since the braid group B_n has a faithful linear representation over a field of characteristic zero (see [17], [18]) then by Corollary 1, the group $\text{Int}(B_n)B_n$ also has a faithful linear representation over the same field. Since $\text{Hol}(B_n)$ is a finite extension of $\text{Int}(B_n)B_n$ then the required statement follows.

Corollary 3. *The holomorph $\text{Hol}(F_2)$ of free group F_2 has a faithful linear representation over a field of characteristic zero.*

Proof. Since the center of F_2 is trivial there is a isomorphism of $\text{Hol}(F_2)$ in the direct production $\text{Aut}(F_2) \times \text{Aut}(F_2)$. To see it we map $\text{Aut}(F_2)$ to diagonal subgroup of this production and F_2 to subgroup $1 \times \text{Int}(F_2)$. Since in $\text{Aut}(F_2)$ the following formulas $\varphi^{-1} \hat{g} \varphi = \hat{g}^{\varphi}$, where $\varphi \in \text{Aut}(F_2)$, $\hat{g} \in \text{Int}(F_2)$, $g \in F_2$, g^{φ} is the image of g under the action φ , are true then the images of F_2 and $\text{Aut}(F_2)$ in $\text{Aut}(F_2) \times \text{Aut}(F_2)$ generate a subgroup which is isomorphic to $\text{Hol}(F_2)$.

Since $\text{Aut}(F_2)$ is linear over a field of characteristic zero (see [20], [21]) then the required assertion follows.

§ 2. On linearity of some HNN–extensions of free group

In this Section we consider HNN–extensions

$$F_{\varphi}(X) = \langle X, t \mid t^{-1} x t = \varphi(x), x \in X \rangle$$

of a free group $F(X)$ by an automorphism $\varphi \in \text{Aut}(F(X))$. We will prove that if the automorphism φ is virtually inner then $F_{\varphi}(X)$ is linear. Note that the conjugacy problem for such groups $F_{\varphi}(X)$ was solved in [13].

The following property holds.

Theorem 2. *Let $F(X)$ be a free group with the a of free generators X , let $\varphi \in \text{Aut } F(X)$ be such that $\varphi^n \in \text{Int } F(X)$ for some positive integer n . If $\sigma : F(X) \longrightarrow \text{GL}_m(P)$ is a faithful linear representation of $F(X)$ by matrices over a field P then the representation τ of $F_\varphi(X)$ defining on generators by the following manner:*

$$t^\tau = \begin{pmatrix} \mathbf{0} & E_m & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & E_m & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & E_m \\ sf^\sigma & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

$$x^\tau = \begin{pmatrix} x^\sigma & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & (\varphi^{-1}(x))^\sigma & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \cdots & (\varphi^{-n+2}(x))^\sigma & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & (\varphi^{-n+1}(x))^\sigma \end{pmatrix},$$

where $f \in F(X)$ is such that $\varphi^n(x) = f^{-1}xf$, $x \in X$, and $s \in P^*$ is an arbitrary element of infinite order, $\mathbf{0}$ and E_m are nullity and unity matrices by $m \times m$ and t^τ and x^τ are matrices by $mn \times mn$ is a faithful linear representation.

Proof. Let

$$F_{\varphi^n}(X) = \langle F(X), t^n \rangle \leq F_\varphi(X).$$

It is obvious that $|F_\varphi(X) : F_{\varphi^n}(X)| = n$, and $F_{\varphi^n}(X)$ is isomorphic to $\mathbb{Z} \times F(X)$. Let us define the representation

$$\bar{\sigma} : F_{\varphi^n}(X) \longrightarrow \langle sE_m, F^\sigma(X) \rangle \leq \text{GL}_m(K),$$

where $(t^n)^{\bar{\sigma}} = sf^\sigma$, $x^{\bar{\sigma}} = x^\sigma$, $x \in X$ and $s \in P^*$ is an element of infinite order.

The representation τ of $F_\varphi(X)$ will construct with the known manner from the representation $\bar{\sigma}$ of subgroup of finite index $F_{\varphi^n}(X)$ (see [1], [19, p. 164]).

If $\{1, t, t^2, \dots, t^{n-1}\}$ is the set of right coset representatives of $F_\varphi(X)$ by subgroup $F_{\varphi^n}(X)$ then for $g \in F_\varphi(X)$ the image g^τ is the matrix of degree mn which consists of blocks of degree m . The block on position (i, j) is equal to $(t^{i-1}gt^{1-j})^{\bar{\sigma}}$ if $t^{i-1}gt^{1-j} \in F_{\varphi^n}(X)$ and equal to a null matrix in the opposite case.

In the matrix t^τ the non-null blocks are the only blocks on the positions $(i, i+1)$, $i = 1, 2, \dots, n-1$. The identical matrix E_m stands on these positions. On the position $(n, 1)$ stands the block $(t^n)^{\bar{\sigma}} = sf^\sigma$. For $x \in X$ in the matrix x^τ the non-null blocks are the only diagonal-blocks and the block on the position (i, i) is equal to $(t^{i-1}xt^{1-i})^{\bar{\sigma}} = (\varphi^{-i+1}(x))^\sigma$. This completes the proof.

The following proposition gives the center of $F_\varphi(X)$.

Proposition 1. *Let*

$$F_\varphi(X) = \langle X, t \mid t^{-1}xt = \varphi(x), \quad x \in X \rangle$$

be an extension of a free group $F(X)$ by a virtually inner automorphism $\varphi \in \text{Aut}(F(X))$. Then the center $Z(F_\varphi(X))$ of $F_\varphi(X)$ is the infinite cyclic group which generated by $t^n w_0$, where n is a minimal natural number for which $\varphi^n \in \text{Int}(F(X))$, and w_0 is defined from the equality $\varphi^n(x) = w_0 x w_0^{-1}$, $x \in X$.

Proof. It is evident that $t^n w_0 \in Z(F_\varphi(X))$ and $(t^n w_0)^k = t^{nk} w_0^k$. Let $h \in Z(F_\varphi(X))$. As element from $F_\varphi(X)$ it can be presented in the form $t^l f$, where $l \in \mathbb{Z}$, $f \in F(X)$. Since h commutes with every $g \in F_\varphi(X)$ then $h^{-1}gh = f^{-1}t^{-l}gt^l f = g$ or $t^{-l}gt^l = f g f^{-1}$. Dividing l with residue on n we get $l = nk + r$, $0 \leq r < n$. Then $t^{-l}gt^l = t^{-nk}t^{-r}gt^r t^{nk} = f g f^{-1}$ or $t^{-r}gt^r = t^{nk} f g f^{-1} t^{-nk} = t^{nk} w_0^k w_0^{-k} f g f^{-1} w_0^k w_0^{-k} t^{-nk} = w_0^{-k} f g f^{-1} w_0^k$. Therefore $r = 0$ and $w_0^{-k} f \in Z(F(X))$. So $t^l f = t^{nk} w_0^k = (t^n w_0)^k \in \langle t^n w_0 \rangle$. The proposition is proven.

§ 3. 2-generated Artin groups

In [22] there was formulated the question on linearity of Artin groups. In this Section we will consider 2-generated Artin groups $A(m)$ which have the following genetic code

$$A(m) = \langle x, y \mid w_m(x, y) = w_m(y, x) \rangle,$$

where $m \geq 3$ and

$$w_m(u, v) = \begin{cases} (uv)^n, & \text{if } m = 2n, \\ (uv)^n u, & \text{if } m = 2n + 1. \end{cases}$$

As it was proved in [13] each 2-generated Artin group $A(m)$ is an extension of a free group of finite rank by some virtually inner automorphism. We will use this result and the results of previous Section to construct some linear representations of $A(m)$ over field of arbitrary characteristic. In particular, we will construct the representation over a ring $\mathbb{Q}_p$ for every prime p , where $\mathbb{Q}_p = \{\frac{x}{p^n} \mid x \in \mathbb{Z}, n \in \mathbb{N} \cup \{0\}\}$. Note, that Squier [14] constructed a faithful linear representation of dimension 2 for the group $A(m)$ over some extension of $\mathbb{Q}$ which has transcendental elements.

Let $\sigma = \sigma(\lambda, \mu) : F_n \longrightarrow \text{GL}_2(\mathbb{Z}[\lambda, \mu])$, $\lambda, \mu \in \mathbb{C}$ be the two-parameter representation of a free group F_n in group of matrices $\text{GL}_2(\mathbb{Z}[\lambda, \mu])$, which is defined on the generators by the rule

$$x_0^\sigma \longmapsto \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix},$$

$$x_i^\sigma \longrightarrow \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}^{-i} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}^i, \quad i = 1, 2, \dots, n-1.$$

It is evident, that this representation is faithful if the group generated by the matrices

$$\begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}$$

is isomorphic to the free group F_2 .

Proposition 2. *Let $A(2n)$, $n \geq 2$, be the 2-generated Artin group. Then the representation*

$$\rho : A(2n) \longrightarrow \mathrm{GL}_{2n}(\mathbb{Z}[\lambda, \mu, s^{\pm 1}]), \quad \lambda, \mu \in \mathbb{C}, \quad s \in \mathbb{C}^*,$$

which is defined on the generators by the rule

$$x^\rho = \mathrm{diag} \left(\begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}, \dots, \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \right), \quad y^\rho = \begin{pmatrix} \mathbf{0} & A & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & A & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \ddots & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & A \\ B & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

where

$$A = \begin{pmatrix} 1 & -\mu \\ 0 & 1 \end{pmatrix}, \quad B = s \begin{pmatrix} 1 & 0 \\ -\lambda & 1 \end{pmatrix} \begin{pmatrix} 1 - \lambda\mu & \mu \\ -\lambda & 1 \end{pmatrix}^{n-1}$$

is a linear representation of $A(2n)$. In addition, if $|\lambda| \geq 2$, $|\mu| \geq 2$ and s is not a root of unity then this representation is faithful.

Proof. We have

$$A(2n) = \langle x_0, x_1, \dots, x_{n-1}, t \mid t^{-1}x_it = \varphi(x_i), \quad i = 0, 1, \dots, n-1 \rangle,$$

where the automorphism φ is defined by the rule

$$\varphi(x_0) = x_0x_1 \dots x_{n-2}x_{n-1}x_{n-2}^{-1} \dots x_1^{-1}x_0^{-1}, \quad \varphi(x_i) = x_{i-1}, \quad i = 1, \dots, n-1.$$

In this case $\varphi^n(x_i) = \Delta x_i \Delta^{-1}$, $\varphi(\Delta) = \Delta$, where $\Delta = x_0x_1 \dots x_{n-1}$ and x_0, t are corresponding to canonical generators of Artin group, i. e.

$$A(2n) = \langle x_0, t \mid (x_0t)^n = (tx_0)^n \rangle,$$

(see [13]).

By Theorem 2 the following matrices are corresponding to generators $x_0, x_1, \dots, x_{n-1}, t$:

$$\begin{aligned} x_0^\tau &= \mathrm{diag}(x_0^\sigma, (\varphi^{-1}(x_0))^\sigma, \dots, (\varphi^{-n+1}(x_0))^\sigma), \\ &\vdots \\ x_{n-1}^\tau &= \mathrm{diag}(x_{n-1}^\sigma, (\varphi^{-1}(x_{n-1}))^\sigma, \dots, (\varphi^{-n+1}(x_{n-1}))^\sigma), \\ t^\tau &= \begin{pmatrix} \mathbf{0} & E_2 & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & E_2 & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \ddots & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & E_2 \\ s(\Delta^{-1})^\sigma & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix}, \end{aligned}$$

where all matrices have degree $2n$, and $s \in \mathbb{C}^*$ is not a root of unity.

If we conjugate t^τ and x_0^τ by the matrix

$$u = \text{diag} \left(E_2, \begin{pmatrix} 1 & -\mu \\ 0 & 1 \end{pmatrix}, \dots, \begin{pmatrix} 1 & -\mu \\ 0 & 1 \end{pmatrix}^{n-1} \right),$$

then we get correspondingly

$$u^{-1}t^\tau u = \begin{pmatrix} \mathbf{0} & A & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & A & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & A \\ B & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad u^{-1}x_0^\tau u = \text{diag} \left(\begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}, \dots, \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \right),$$

where

$$A = \begin{pmatrix} 1 & -\mu \\ 0 & 1 \end{pmatrix}, \quad B = s \begin{pmatrix} 1 & 0 \\ -\lambda & 1 \end{pmatrix} \begin{pmatrix} 1 - \lambda\mu & \mu \\ -\lambda & 1 \end{pmatrix}^{n-1}.$$

Let us define a representation ρ taking $x^\rho = u^{-1}x_0^\tau u$, $y^\rho = u^{-1}t^\tau u$. The elements $x = x_0$ and $y = t$ are canonical generators of Artin group $A(2n)$. If in the representation $\sigma = \sigma(\lambda, \mu)$ we take $|\lambda| \geq 2$, $|\mu| \geq 2$, and s is a non-zero complex number not equal to a root of unity, then ρ is faithful representation. If we take $\lambda = \mu = 2$, $s = p$ is a prime number then we get a faithful representation of $A(2n)$, $n \geq 2$ by matrices of degree $2n$ over the ring $\mathbb{Q}_p$. This completes the proof.

In the case when $m = 2n + 1$, $n \geq 1$ is odd number, the 2-generated Artin group $A(m)$ is isomorphic to

$$F_{2n+1}(\psi) = \langle x_0, x_1, \dots, x_{2n-1}, t \mid t^{-1}x_it = \psi(x_i), \quad i = 0, 1, \dots, 2n-1 \rangle,$$

where the automorphism ψ is given by

$$\psi(x_0) = x_0x_2 \dots x_{2n-4}x_{2n-2}x_{2n-1}^{-1} \dots x_3^{-1}x_1^{-1}, \quad \psi(x_i) = x_{i-1}, \quad i = 1, \dots, 2n-1.$$

In this case $\psi^{2(2n+1)}(x_i) = \Sigma x_i \Sigma^{-1}$, where

$$\Sigma = x_0x_2 \dots x_{2n-2}(x_0x_1 \dots x_{2n-1})^{-1}x_1x_3 \dots x_{2n-1}, \quad \psi(\Sigma) = \Sigma$$

(see [13]). Note that the inverse automorphism ψ^{-1} is given by

$$\psi^{-i}(x_0) = x_i, \quad i = 1, 2, \dots, 2n-1,$$

$$\psi^{-i}(x_0) = \Sigma^{-1}\psi^{4n+2-i}(x_0)\Sigma, \quad i = 2n, 2n+1, \dots, 4n+1.$$

The canonical generators of $A(2n+1)$ are $x = t$ and $y = x_0t$, i. e.

$$A(2n+1) = \langle t, x_0t \mid (tx_0t)^nt = (x_0t^2)^nx_0t \rangle.$$

Proposition 3. *Let $A(2n + 1)$, $n \geq 1$ be the 2-generated Artin group with odd number of generators. Then the map*

$$\rho : A(2n + 1) \longrightarrow \mathrm{GL}_{4(2n+1)}(\mathbb{Z}[\lambda, \mu, s^{\pm 1}]), \quad \lambda, \mu \in \mathbb{C}, \quad s \in \mathbb{C}^*,$$

which is defined on the generators

$$x^\rho = \begin{pmatrix} \mathbf{0} & E_2 & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & E_2 & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & E_2 \\ s(\Sigma^{-1})^\sigma & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

$$y^\rho = \begin{pmatrix} \mathbf{0} & x_0^\sigma & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & (\psi^{-1}(x_0))^\sigma & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & (\psi^{-4n}(x_0))^\sigma \\ s(\Sigma^{-1}\psi(x_0))^\sigma & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

where $\sigma = \sigma(\lambda, \mu)$ is the representation of a free 2-generated group, is the linear representation of $A(2n + 1)$. In this case, if $|\lambda| \geq 2$, $|\mu| \geq 2$ and s is not a root of unity then this representation is faithful.

Proof. By Theorem 2 we get for elements $x_0, x_1, \dots, x_{2n-1}, t$ the following matrices

$$x_0^\tau = \mathrm{diag}(x_0^\sigma, (\psi^{-1}(x_0))^\sigma, \dots, (\psi^{-(4n+1)}(x_0))^\sigma),$$

$$\vdots$$

$$x_{2n-1}^\tau = \mathrm{diag}(x_{2n-1}^\sigma, (\psi^{-1}(x_{2n-1}))^\sigma, \dots, (\psi^{-(4n+1)}(x_{2n-1}))^\sigma),$$

$$t^\tau = \begin{pmatrix} \mathbf{0} & E_2 & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & E_2 & \cdots & \mathbf{0} & \mathbf{0} \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & E_2 \\ s(\Sigma^{-1})^\sigma & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

where all matrices have degree $4n + 2$, and $s \in \mathbb{C}^*$ is not a root of unity. The map $x \longmapsto t^\tau$, $y \longmapsto x_0^\tau t^\tau$ define of desired representation.

In view of Theorem 2, if the representation σ is faithful then τ is the faithful representation of $F_{2n+1}(\psi)$. In this case, the matrices t^τ , $x_0^\tau t^\tau$ are canonical generators of $A(2n + 1)$. This completes the proof.

Note, that for $\lambda = \mu = 2$ and $s = p$ is a prime number, we get a representation of $A(2n + 1)$ by matrices of degree $4n + 2$ over the ring $\mathbb{Q}_p$.

Since 2-generated Artin group $A(m)$ is a split extension of a free group of finite rank by a virtually inner automorphism, then using of Lemma 4 we find the center of $A(m)$.

Corollary. *The center of 2-generated Artin group $A(m)$ is infinite cyclic group. In this case*

- a) *if $m = 2n$ then the center $Z(A(m))$ is generated by $(xy)^n$,*
- b) *if $m = 2n + 1$ then the center $Z(A(m))$ is generated by $((xy)^n x)^2$.*

Proof. a) We have the isomorphism

$$A(2n) \simeq \langle x_0, x_1, \dots, x_{n-1}, t \mid t^{-1}x_it = \varphi(x_i), \quad i = 0, 1, \dots, n-1 \rangle$$

and $\varphi^n(x_i) = \Delta x_i \Delta^{-1}$, $\Delta = x_0 x_1 \dots x_{n-1}$, $\varphi(\Delta) = \Delta$. From the definition of φ it follows that n is the least positive power for which the automorphism φ^n is inner. Since for this isomorphism to $t \in F_\varphi(X)$ corresponds the element $y \in A(2n)$, and to $x_0 \in F_\varphi(X)$ corresponds the element $x \in A(2n)$ then to $(x_0 t)^n$ corresponds the element $(xy)^n$. To use Proposition 1 we note that

$$(x_0 t)^n = x_0 x_1 \dots x_{n-1} t^n = \Delta t^n = t^n \Delta.$$

b) We have the isomorphism

$$A(2n+1) \simeq \langle x_0, x_1, \dots, x_{2n-1}, t \mid t^{-1}x_it = \psi(x_i), \quad i = 0, 1, \dots, 2n-1 \rangle$$

and $\psi^{2(2n+1)}(x_i) = \Sigma x_i \Sigma^{-1}$, $\Sigma = x_0 x_2 \dots x_{2n-2} (x_0 x_1 \dots x_{2n-1})^{-1} x_1 x_3 \dots x_{2n-1}$, $\psi \Sigma = \Sigma$. Since the elements $t = x$ and $x_0 t = y$ are canonical generators of $A(2n+1)$ then element $(tx_0 t)^n t$ maps to $(xy)^n x$. Further, since $((tx_0 t)^n t)^2 = t^{4n+2} \Sigma$ and $4n+2$ is a least positive power in which the automorphism ψ is inner then from proposition 1 follows desired statement.

Note that the center of Artin groups was found in the paper of E. Briskorn and K. Saito [23].

We will construct a series of representations for $A(3)$. This group is isomorphic to the braid group on 3 strings B_3 .

Using Lemma 3 we take

$$X_0 = \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix}, \quad X_1 = \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}, \quad \lambda, \mu \in \mathbb{C}.$$

Then the following matrices

$$T = \begin{pmatrix} \mathbf{0} & E_2 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & E_2 & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 \\ s\Sigma^{-1} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

DT , where $D = \text{diag}(X_0, X_1, \psi^{-2}(X_0), \psi^{-3}(X_0), \psi^{-4}(X_0), \psi^{-5}(X_0))$ correspond to generators. If we conjugate these matrices by the matrix

$$U = \text{diag}(E_2, E_2, \Sigma^{-1}, \Sigma^{-1}, \Sigma^{-1}, \Sigma^{-1}),$$

then we get

$$U^{-1}TU = \begin{pmatrix} \mathbf{0} & E_2 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \Sigma^{-1} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & E_2 \\ sE_2 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

$$U^{-1}DTU = \begin{pmatrix} \mathbf{0} & X_0 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & X_1\Sigma^{-1} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \psi^4(X_0) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \psi^3(X_0) & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \psi^2(X_0) \\ s\psi(X_0) & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix},$$

We take these matrices as canonical generators X and Y of $A(3)$. It is easily show that the following relation is true: $XYX = YXY$.

This representation will be faithful if the group generated by X_0 and X_1 is isomorphic to the free group F_2 . Using direct calculations we find the matrices

$$\Sigma^{-1} = \begin{pmatrix} 1 - \lambda\mu + \lambda^2\mu^2 & -\lambda\mu^2 \\ -\lambda^2\mu & 1 + \lambda\mu \end{pmatrix}, \quad \psi(X_0) = \begin{pmatrix} 1 & -\mu \\ \lambda & 1 - \lambda\mu \end{pmatrix},$$

$$\psi^2(X_0) = \begin{pmatrix} 1 + \lambda\mu & -\mu \\ \lambda^2\mu & 1 - \lambda\mu \end{pmatrix}, \quad \psi^3(X_0) = \begin{pmatrix} 1 + \lambda\mu - \lambda^2\mu^2 & \lambda\mu^2 \\ -\lambda(1 - \lambda\mu)^2 & 1 - \lambda\mu + \lambda^2\mu^2 \end{pmatrix},$$

$$\psi^4(X_0) = \begin{pmatrix} 1 - 2\lambda^2\mu^2 & \mu(1 + 2\lambda\mu) \\ \lambda(1 + \lambda\mu - 2\lambda^2\mu^2) & 1 - \lambda\mu + 2\lambda^2\mu^2 \end{pmatrix},$$

where $s \in \mathbb{C}^*$ and is not a root of unity.

If in Theorem 2 for $\bar{\sigma}$ we take the representation

$$(t^n)^{\bar{\sigma}} = \begin{pmatrix} T(s) & 0 \\ 0 & f^\sigma \end{pmatrix}, \quad x^{\bar{\sigma}} = \begin{pmatrix} E_2 & 0 \\ 0 & x^\sigma \end{pmatrix}, \quad T(s) = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix},$$

where $\sigma : F(X) \longrightarrow \text{SL}_2(\mathbb{Z})$, $s \in \mathbb{Z} \setminus \{0\}$ then we get the representation of $F_\varphi(X)$ in $\text{SL}_{4n}(\mathbb{Z})$. In particular, for the braid group B_3 which is isomorphic to $A(3)$ we can take $\lambda = \mu = 2$, $s \in \mathbb{Z} \setminus \{0\}$ and get the representation $\bar{\sigma}$ (see Proposition 3) which is faithful representation by integer valued matrices of dimension 24.

REFERENCES

1. A. I. Malcev, On isomorphic matrix representations of infinite groups. (Russian; with English summary), Mat. Sb., 8, № 3 (1940), 405–422.
2. Yu. I. Merzljakov, Linear groups. (Russian, English) J. Sov. Math., 1, (1973) 571–593; translation from Itogi Nauki, Ser. Mat., Algebra, Topologija, Geometrija 1970, 75–110 (1971).
3. A. A. Lubotzky, Group-theoretic characterization of linear groups, J. Algebra, 113 (1988), 207–214.
4. V. L. Nisnevich, Über Gruppen die durch Matrizen über einem kommutativen Feld isomorph darstellbar sind (Russian, German summary), Mat. Sb., 8, № 3 (1940), 395–403.
5. B. A. F. Wehrfritz, Generalized free products of linear groups, Proc. London Math. Soc., 27, (1973), 402–424.
6. Yu. E. Vapne, On the representability of a direct interlacing of groups by matrices. (Russian, English) Math. Notes 7, (1970), 110–114; translation from Mat. Zametki 7, (1970), 181–189.
7. Yu. E. Vapne, Matrix representability of nilpotent products of groups. (Russian) Math. Notes, 14, № 3 (1973), 383–394.
8. O. V. Bryukhanov, Matrix representability and structure of groups, Algebra and model theory – 4, Novosibirsk State Technical University, Novosibirsk, 2003, 15–29.
9. R. T. Volvachev, Linear representability of the groups $G_{n,k,l} = \langle a, t \mid t^{-1}a^kt = a^l \rangle$. (Russian. English summary), Fundam. Prikl. Mat., 4, № 4 (1998), 1415–1418.
10. Yu. I. Merzljakov, Matrix representation of automorphisms, extensions, and solvable groups, Algebra and Logic, 7, (1970), 169–192.
11. Yu. I. Merzljakov, Rational groups (Russian), M.: Nauka, 1980.
12. V. Bardakov, L. Bokut and A. Vesnin, Twisted conjugacy in free groups and Makanin’s question, Southeast Asian Bulletin of Math., 29, № 2 (2005), 209–226. (See also arXiv:math.GR/0401349.)
13. V. Bardakov, L. Bokut and A. Vesnin, On the conjugacy problem for cyclic extensions of free groups, Comm. Algebra, 33, № 6 (2005), 1979–1996. (See also arXiv:math.GR/0402060.)
14. C. C. Squier, Matrix representations of Artin groups, Proc. Amer. Math. Soc., 103, № 1 (1988), 49–53.
15. B. A. F. Wehrfritz, Infinite linear groups, Berlin, 1973.

16. J. L. Dyer, E. K. Grossman, The automorphism groups of the braid groups, *Amer. J. Math.*, 103, № 6 (1981), 1151–1169.
17. S. Bigelow, Braid groups are linear, *J. Amer. Math. Soc.*, 14, № 2 (2001), 471–486.
18. D. Krammer, Braid groups are linear, *Annals of. Math.*, 155, № 1 (2002), 131–156.
19. M. I. Kargapolov, Yu. I. Merzljakov, *Fundamentals of the theory of groups*. Transl. from the 2nd Russian ed. by Robert G. Burns. (Russian, English) *Graduate Texts in Mathematics*. 62. New York-Heidelberg-Berlin: Springer-Verlag. XVII, 203 p. (1979).
20. Bardakov V. G. Linear representations of the group of conjugating automorphisms and the braid groups of some manifolds, *Sibirsk. Mat. Zh.*, 46 N 1 (2005), 17–31.
 Линейные представления группы сопрягающих автоморфизмов и групп кос некоторых многообразий, *Сиб. матем. журн.*, 46, № 1 (2005), 17–31.
21. Dyer J. L., Formanek E. and Grossman E. K. On the linearity of automorphism groups of free groups, *Arch. Math.*, 38, N 5 (1982), 404–409.
22. T. Hsu, D. T. Wise, On linear and residual properties of graph products, *Michigan Math. J.*, 46, № 2 (1999), 251–259.
23. E. Brieskorn, K. Saito. *Artin-Gruppen und Coxeter-Gruppen*. (Russian) *Matematika*, Moskva 18, № 6 (1974), 56–79.

Sobolev Institute of Mathematics, Novosibirsk 630090,
 Russia, e-mail: bardakov@math.nsc.ru